

HIFEM[®] FOR FEMALE SEXUAL DYSFUNCTION

THE USE OF HIFEM TECHNOLOGY IN THE TREATMENT OF PELVIC FLOOR MUSCLES AS A CAUSE OF FEMALE SEXUAL DYSFUNCTION: A MULTI-CENTER PILOT STUDY

Timothy Charles Hlavinka MD¹, Pavel Turčan MD², Alexander Bader MD, FAAOCG, FAAC³

1. The Urology Place, InVita Clinics, San Antonio USA


2. Center Mediol, Olomouc, Czech Republic

3. Bader Medical Institute of London, Marylebone, London, United Kingdom

Published in Journal of Women's Health Care: March 9, 2019

HIGHLIGHTS

- Average FSFI score **increased** on average by **53%** after six treatments.
- **Improvement** rate was **maintained** during 3-month follow-up.
- Most prominent improvement was seen in sexual satisfaction (**+76%**), desire (**+76%**) and orgasm (**+60%**) domains.
- HIFEM procedure significantly improved female sexual function through strengthening of pelvic floor muscles.



Changes in FSFI full score in treated patients. Blue rectangles contain 50% of data; whiskers indicate variability of the results. Improvement after the six procedures (+53%) was maintained at 3 months (+51%).